

INBRED CORN LINE G3001**FIELD OF THE INVENTION**

5 This invention is in the field of corn breeding, specifically relating to an inbred corn line designated G3001. This invention also is in the field of hybrid maize production employing the present inbred.

BACKGROUND OF THE INVENTION

10 The original maize plant was indigenous to the Western Hemisphere. The plants were weed-like and only through the efforts of early breeders were cultivated crop species developed. The crop cultivated by early breeders, like the crop today, could be wind pollinated. The physical traits of maize are such that
15 wind pollination results in self-pollination or cross-pollination between plants. Each plant has a separate male and female flower that contributes to pollination, the tassel and ear, respectively. Natural pollination occurs when wind transfers pollen from tassel to the silks on the corn ears. This type of pollination has contributed to the wide variation of maize varieties present in the Western Hemisphere.

20 The development of a planned breeding program for maize only occurred in the last century. A large part of the development of the maize product into a profitable agricultural crop was due to the work done by land grant colleges. Originally, maize was an open pollinated variety having heterogeneous genotypes.
25 The maize farmer selected uniform ears from the yield of these genotypes and preserved them for planting the next season. The result was a field of maize plants that were segregating for a variety of traits. This type of maize selection led to, at most, incremental increases in seed yield.

30 Large increases in seed yield were due to the work done by land grant

colleges that resulted in the development of numerous hybrid corn varieties in planned breeding programs. Hybrids were developed by selecting corn lines and selfing these lines for several generations to develop homozygous pure inbred lines. One selected inbred line was crossed with another selected inbred line to produce hybrid progeny (F1). Although hybrids, due to heterosis, are robust and vigorous plants. Inbreds on the other hand are mostly homozygous. This homozygosity renders the inbred lines less vigorous. Inbred seed can be difficult to produce since the inbreeding process in corn lines decreases the vigor. However, when two inbred lines are crossed, the hybrid plant evidences greatly increased vigor and seed yield compared to open pollinated, segregating maize plants. An important consequence of the homozygosity and the homogeneity of the inbred maize lines is that all hybrid seed produced from any cross of two such elite lines will be the same hybrid seed and make the same hybrid plant. Thus the use of inbreds makes hybrid seed which can be reproduced readily. The hybrid plant in contrast does not produce hybrid seed that is readily reproducible. The seed on a hybrid plant is segregating for traits.

The ultimate objective of the commercial maize seed companies is to produce high yielding, agronomically sound plants that perform well in certain regions or areas of the Corn Belt. To produce these types of hybrids, the companies must develop inbreds, which carry needed traits into the hybrid combination. Hybrids are not often uniformly adapted for the entire Corn Belt, but most often are specifically adapted for regions of the Corn Belt. Northern regions of the Corn Belt require shorter season hybrids than do southern regions of the Corn Belt. Hybrids that grow well in Colorado and Nebraska soils may not flourish in richer Illinois and Iowa soil. Thus, a variety of major agronomic traits are important in hybrid combination for the various Corn Belt regions, and have an impact on hybrid performance.

Inbred line development and hybrid testing have been emphasized in the past half-century in commercial maize production as a means to increase hybrid performance. Inbred development is usually done by pedigree selection. Pedigree selection can be selection in an F_2 population produced from a planned cross of two genotypes (often elite inbred lines), or selection of progeny of synthetic varieties, open pollinated, composite, or backcrossed populations. This type of selection is effective for highly inheritable traits, but other traits, for example, yield requires replicated test crosses at a variety of stages for accurate selection.

Maize breeders select for a variety of traits in inbreds that impact hybrid performance along with selecting for acceptable parental traits. Such traits include: yield potential in hybrid combination; dry down; maturity; grain moisture at harvest; greensnap; resistance to root lodging; resistance to stalk lodging; grain quality; disease and insect resistance; ear and plant height. Additionally, Hybrid performance will differ in different soil types such as low levels of organic matter, clay, sand, black, high pH, low pH; or in different environments such as wet environments, drought environments, and no tillage conditions. These traits appear to be governed by a complex genetic system that makes selection and breeding of an inbred line extremely difficult. Even if an inbred in hybrid combination has excellent yield (a desired characteristic), it may not be useful because it fails to have acceptable parental traits such as seed yield, seed size, pollen production, good silks, plant height, etc.

To illustrate the difficulty of breeding and developing inbred lines, the following example is given. Two inbreds compared for similarity of 29 traits differed significantly for 18 traits between the two lines. If 18 simply inherited single gene traits were polymorphic with gene frequencies of 0.5 in the parental lines, and assuming independent segregation (as would essentially be the case if

each trait resided on a different chromosome arm), then the specific combination of these traits as embodied in an inbred would only be expected to become fixed at a rate of one in 262,144 possible homozygous genetic combinations. Selection of the specific inbred combination is also influenced by the specific selection environment on many of these 18 traits which makes the probability of obtaining this one inbred even more remote. In addition, most traits in the corn genome are regrettably not single dominant genes but are multi-genetic with additive gene action not dominant gene action. Thus, the general procedure of producing a non segregating F_1 generation and self pollinating to produce a F_2 generation that segregates for traits and selecting progeny with the visual traits desired does not easily lead to a useful inbred. Great care and breeder expertise must be used in selection of breeding material to continue to increase yield and the agronomics of inbreds and resultant commercial hybrids.

Certain regions of the Corn Belt have specific difficulties that other regions may not have. Thus the hybrids developed from the inbreds have to have traits that overcome or at least minimize these regional growing problems. Examples of these problems include in the eastern corn belt Gray Leaf Spot, in the north cool temperatures during seedling emergence, in the Nebraska region CLN (corn Lethal necrosis and in the west soil that has excessively high pH levels. The industry often targets inbreds that address these issues specifically forming niche products. However, the aim of most large seed producers is to provide a number of traits to each inbred so that the corresponding hybrid can be useful in a broader regions of the Corn Belt. The new biotechnology techniques such as Microsatellites, RFLPs, RAPDs and the like have provided breeders with additional tools to accomplish these goals.

SUMMARY OF THE INVENTION

The present invention relates to an inbred corn line G3001. Specifically, this invention relates to plants and seeds of this line. Additionally, this relates to a method of producing from this inbred, hybrid seed corn and hybrid plants with seeds from such hybrid seed. More particularly, this invention relates to the unique combination of traits that combine in corn line G3001.

Generally then, broadly the present invention includes an inbred corn seed designated G3001. This seed produces a corn plant.

The invention also includes the tissue culture of regenerable cells of G3001 wherein the cells of the tissue culture regenerates plants capable of expressing the genotype of G3001. The tissue culture is selected from the group consisting of leaves, pollen, embryos, roots, root tips, guard cells, ovule, seeds, anthers, silk, flowers, kernels, ears, cobs, husks and stalks, cells and protoplasts thereof. The corn plant regenerated from G3001 or any part thereof is included in the present invention. The present invention includes regenerated corn plants that are capable of expressing G3001's genotype, phenotype or mutants or variants thereof.

The invention extends to hybrid seed produced by planting, in pollinating proximity which includes using preserved maize pollen as explained in U.S. Patent 5,596,838 to Greaves , seeds of corn inbred lines G3001 and another inbred line if preserved pollen is not used; cultivating corn plants resulting from said planting; preventing pollen production by the plants of one of the inbred lines if two are employed; allowing cross pollination to occur between said inbred lines; and harvesting seeds produced on plants of the selected inbred. The hybrid seed produced by hybrid combination of plants of inbred corn seed designated G3001

and plants of another inbred line are apart of the present invention. This inventions scope covers hybrid plants and the plant parts including the grain and pollen grown from this hybrid seed.

5 The invention further includes a method of hybrid F1 production. A first generation (F1) hybrid corn plant produced by the process of planting seeds of corn inbred line G3001; cultivating corn plants resulting from said planting; permitting pollen from another inbred line to cross pollinate inbred line G3001; harvesting seeds produced on plants of the inbred; and growing a harvested seed
10 are part of the method of this invention.

Likewise included is a first generation (F1) hybrid corn plant produced by the process of planting seeds of corn inbred line G3001; cultivating corn plants resulting from said planting; permitting pollen from inbred line G3001 to cross
15 pollinate another inbred line; harvesting seeds produced on plants of the inbred; and growing a plant from such a harvested seed.

The inbred corn line G3001 and at least one transgenic gene adapted to give G3001 additional and/or altered phenotypic traits are within the scope of the invention. Such transgenes are usually associated with regulatory elements
20 (promoters, enhancers, terminators and the like). Presently, transgenes provide the invention with traits such as insect resistance, herbicide resistance, disease resistance increased or decreased starch or sugars or oils, increased or decreased life cycle or other altered trait.

25 The present invention includes inbred corn line G3001 and at least one transgenic gene adapted to give G3001 modified starch traits. Furthermore this invention includes the inbred corn line G3001 and at least one mutant gene adapted to give modified starch, acid or oil traits. The present invention includes

the inbred corn line G3001 and at least one transgenic gene selected from the group consisting of: bacillus thuringiensis, the bar or pat gene encoding Phosphinothricin acetyl Transferase, Gdha gene, EPSP synthase gene, low phytic acid producing gene, and zein. The inbred corn line G3001 and at least one transgenic gene useful as a selectable marker or a screenable marker are covered by the present invention.

A tissue culture of the regenerable cells of hybrid plants produced with use of G3001 genetic material is covered by this invention. A tissue culture of the regenerable cells of the corn plant produced by the method described above are also included.

DEFINITIONS

In the description and examples, which follow, a number of terms are used. In order to provide a clear and consistent understanding of the specifications and claims, including the scope to be given such terms, the following definitions are provided.

BL MOIST

The moisture percentage of the grain at black layer, i.e., when 50% of the plants per plot have reached physiological maturity.

COLD GERM

Cold Germ is a measurement of seed germination under cold soil conditions. Data is reported as percent of seed germinating.

ECB

European corn borer is a maize eating insect. ECBI is the first brood generation of

European corn borers. ECBII is the second generation of European corn borers.

EMERGE (EMG)

The number of emerged plants per plot (planted at the same seedling rate) collected when plants have two fully developed leaves.

GI

This is a selection index that provides a single quantitative measure of the worth of a hybrid based on four traits. FI is a very similar index which weights yield less than GI. In GI yield is the primary trait contributing to index values. The GI value is calculated by combining stalk lodging, root lodging, yield and dropped ears according to the attached mathematical formula:

$$GI = 100 + 0.5 (YLD) - 0.9 (\%STALK LODGE) - 0.9 (\%ROOT LODGE) - 2.7 (\%DROPPED EAR)$$

GLS

Gray Leaf Spot (*Cercospora Zeae*) disease rating. This is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.*

GW

Goss' Wilt (*Corynebacterium nebraskense*). This is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.*

HEATP10

The number of Growing Degree Units (GDU's) or heat units required for an inbred line or hybrid to have approximately 10 percent of the plants shedding pollen. This trait is measured from the time of planting. Growing Degree Units are calculated by the Barger Method where the GDU's for a 24 hour period are:

$$\text{GDU} = \frac{(\text{Max Temp } (^{\circ}\text{F}) + \text{Min Temp } (^{\circ}\text{F})) - 50}{2}$$

The highest maximum temperature used is 86°F and the lowest minimum temperature used is 50°F. For each inbred or hybrid it takes a certain number of GDU's to reach various stages of plant development.

HEATBL

The number of GDU's after planting when approximately 50 percent of the inbred or hybrid plants in a plot have grain that has reached physiological maturity (black layer).

HEATPEEK

The number of GDU's after planting of an inbred when approximately 50 percent of the plants show visible tassel extension.

HEATP50 or HTP50

The number of GDU's required for an inbred or hybrid to have approximately 50 percent of the plants shedding pollen. Growing Degree Units are calculated by the

Barger Method as shown in the HEATP10 definition.

HEATP90

5 The number of GDU's accumulated from planting when the last 100 percent of plants in an inbred or hybrid are still shedding enough viable pollen for pollination to occur. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

10 **HEATS10**

The number of GDU's required for an inbred or hybrid to have approximately 10 percent of the plants with silk emergence of at least 0.5 inches. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

15 **HEATS50 or HTS50**

The number of GDU's required for an inbred or hybrid to have approximately 50 percent of the plants with silk emergence of at least 0.5 inches. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

20 **HEATS90**

The number of GDU's required for an inbred or hybrid to have approximately 90 percent of the plants with silk emergence of at least 0.5 inches. Growing Degree Units are calculated by the Barger Method as shown in the HEATP10 definition.

25 **MDMV_A**

Maize Dwarf Mosaic Virus strain A. The corn is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.*

MDMV_B

Maize Dwarf Mosaic Virus strain B. This is rated on a 1-9 scale with a "1" being very susceptible and a "9" being very resistant.*

MOISTURE

The average percentage grain moisture of an inbred or hybrid at harvest time.

NLB

Northern Leaf Blight (*Exserohilum turcicum*) disease rating. This is rated on a 1-9 scale with a "1" being very susceptible, and a "9" being very resistant.*

PCT TILLER

The total number of tillers per plot divided by the total number of plants per plot.

PLANT

This term includes plant cells, plant protoplasts, plant cell tissue cultures from which corn plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as embryos, pollen, flowers, kernels, ears, cobs, leaves, husks, stalks, roots, root tips, anthers, silk and the like., and this term also includes any transgenic DNA or (RNA) or portion thereof that have been introduced into the plant by whatever method.

PLANT HEIGHT (PLTHT) (PHT)

The distance in centimeters from ground level to the base of the tassel peduncle.

5

PLANT INTEGRITY (PLTINT) or (INT)

The level of plant integrity on a scale of 1-9 with 9 evidencing the trait most strongly: 1-2.9 ratings are low plant integrity, 3-5.9 ratings are intermediate plant integrity, and 6-9 ratings are strongly evidencing plant integrity.

10

POPULATION (POP)

The plant population.

RM

Predicted relative maturity based on the moisture percentage of the grain at harvest. This rating is based on known set of checks and utilizes standard linear regression analyses and is referred to as the Minnesota Relative Maturity Rating System.

20

SHED

The volume of pollen shed by the male flower rated on a 1-9 scale where a "1" is a very light pollen shedder, a "4.5" is a moderate shedder, and a "9" is a very heavy shedder.

25

SLB

Southern Leaf Blight (*Bipolaris maydis*) disease rating. This is rated on a 1-9 scale

with a "1" being very susceptible, and a "9" being very resistant.*

STAYGREEN (SGN)

5 The level of staygreen of the plant on a scale of 1-9 with 9 evidencing the trait most strongly: 1-2.9 ratings are low staygreen, 3-5.9 ratings are intermediate staygreen, and 6-9 ratings are strongly evidencing staygreen.

TWT

10 The measure of the weight of grain in pounds for a one bushel volume adjusted for percent grain moisture.

VIGOR (VIG)

15 Visual rating of 1 to 9 made 2-3 weeks post-emergence where a "1" indicates very poor early plant development, and a "9" indicates superior plant development.

WARM GERM

20 A measurement of seed germination under ideal (warm, moist) conditions. Data is reported as percent of seeds germinating.

YIELD (YLD)

25 Actual yield of grain at harvest adjusted to 15.5% moisture. Measurements are reported in bushels per acre.

% DROPPED EARS (DE)

The number of plants per plot, which dropped their primary ear, divided by the total number of plants per plot.

5

% LRG FLAT

Percentage by weight of shelled corn that passes through a 26/64 inch round screen and a 14/64 inch slot screen, but does not pass through a screen with 20.5/64 inch round openings.

10

% LRG ROUND

Percentage by weight of shelled corn that passes through a 26/64 inch round screen, but does not pass through a 14/64 inch slot screen or a screen with 20.5/64 inch round openings.

15

% MED FLAT

Percentage by weight of shelled corn that passes through a 20.5/64 inch round screen and a 13/64 inch slotted screen, but does not pass through a screen with 17/64 inch round openings.

20

% MED ROUND

Percentage by weight of shelled corn that passes through a 20.5/64 inch round screen, but does not pass through a 13/64 inch slot screen or a screen with 17/64 inch round openings.

25

% SML FLAT

Percentage by weight of shelled corn that passes through a 17/64 inch round screen and a 12/64 inch slotted screen, but does not pass through a screen with 15/64 inch round openings.

% SML ROUND

Percentage by weight of shelled corn that passes through a 17/64 inch round screen, but does not pass through a 12/64 inch slotted screen or a screen with 15/64 inch round openings.

% ROOT LODGE (RL)

Percentage of plants per plot leaning more than 30 degrees from vertical divided by total plants per plot.

% STALK LODGE (SL)

Percentage of plants per plot with the stalk broken below the primary ear node divided by the total plants per plot.

*Resistant - on a scale of 1-9 with 9 evidencing the trait most strongly: 1-2.9 ratings are susceptible, 3-5.9 ratings are intermediate, and 6-9 ratings are resistant.

DETAILED DESCRIPTION OF THE INVENTION

G3001 can be used as a female line. This inbred is useful as a female because it has nice seed producing traits. Good yields with a large percentage of the seeds being medium sized flat or rounds. In hybrid combination this line yields nice 108 day hybrids. The inbred carries a slightly better than average plant integrity into late season.

The inbred has shown uniformity and stability within the limits of environmental influence for all the traits as described in the Variety Description Information (Table 1) that follows.

The inbred has been self-pollinated for a sufficient number of generations to give inbred uniformity. During plant selection in each generation, the uniformity of plant type was selected to ensure homozygosity and phenotypic stability. The line has been increased in isolated farmland environments with data on uniformity and agronomic traits being observed to assure uniformity and stability. No variant traits have been observed or are expected in G3001.

The best method of producing the invention, G3001 which is substantially homozygous, is by planting the seed of G3001 which is substantially homozygous and self-pollinating or sib pollinating the resultant plant in an isolated environment, and harvesting the resultant seed.

TABLE 1

G3001

VARIETY DESCRIPTION INFORMATION

- #1 Type: Dent
#2 This inbred has RM of 107.

#3 Plant Traits

Plant Height 76 in.
Ear Height 29 in.
Tillers (Rating) 5
Leaf Color MEDIUM GREEN
Brace Root Color RED
Silk Color RED
Shoots at Flowering BALD

#4 Tassel Traits

Glume Color GREEN/PURPLE
Glume Ring Color RED
Anther Color YELLOW

#5 Ear and Kernel Traits

Cob Color RED
Kernel Crown Color LIGHT YELLOW
Kernel Body Color DARK YELLOW

6 DISEASE RESISTANCE IN INBRED G3001

Eye Spot = 5.1

Gray Leaf Spot = 4.00

Gross wilt = 2.5

Northern leaf blight = 6.0

#7 INSECT RESISTANCE IN INBRED

ECB1 = 3.75

ECB2 = 5.31530

Ear rate = 3.3125

#8 The comparable inbred to G3001 is ZS0560, an inbred having a number of similarities. ZS0560 is an inbred which has been or is presently in a number of commercial hybrids that are in a similar region of adaptation as most of the hybrids formed with G3001.

The Munsell code is a reference book of color, which is known and used in the industry and by persons with ordinary skill in the art of plant breeding.

The purity and homozygosity of inbred G3001 is constantly being tracked using isozyme genotypes as shown in Table 2.

Isozyme Genotypes for G3001

Isozyme data were generated for inbred corn line G3001 according to procedures known and published in the art. The data in Table 2 gives the electrophoresis data on G3001 as compared to its two parents.

TABLE 2
ELECTROPHORESIS RESULTS FOR G3001

	INBRED	ACP1	ACP4	ADH	MDH1	MDH2	PGD1	PGD2	PHI	PGM	IDH
5	G3001	33	55	22	22	11	22	22	22	22	22

Inbred and Hybrid Performance of G3001

10 The traits and characteristics of inbred corn line G3001 are listed to compare with other inbreds and/or in hybrid combination. The G3001 data shows the characteristics and traits of importance, giving a snapshot of G3001 in these specific environments.

15 Table 3A shows a comparison between G3001 and a comparable inbred ZS0560. G3001 has excellent vigor compared to inbred ZS0560. The two inbreds show significant differences in ear height, plant height, and across all Heat measurements for pollination and silking. G3001 has significantly higher yield at harvest than does ZS0560 but has slightly higher moisture. G3001 reaches heat
20 peek with significantly less heat units than does ZS0560. The present invention is slightly less full season than is ZS0560. G3001 has significantly less medium round seeds and significantly more medium flat seeds than does ZS0560.

TABLE 3A G3001

YEAR	INBRED	VIGOR	EMERGE	PCT TILLER	PLANT HEIGHT	EAR HEIGHT	SHED	EAR QUALITY	PCT BARREN
OVERALL	G3001	6.4	84.9		174.7	63.3	3.7		
	ZS0560	5.5	84.0		165.2	68.9	4.3		
	# EXPTS	18	31		24	22	23		
	DIFF	0.9	0.8		9.5	5.6	0.6		
	PROB	0.001***	0.547		0.002***	0.016**	0.016		

YEAR	INBRED	HEATP10	HEATP50	HEATP90	HEATS10	HEATS50	HEATS90
OVERALL	G3001	1412	1456	1538	1407	1447	1476
	ZS0560	1464	1503	1606	1462	1502	1524
	# EXPTS	34	34	16	34	34	21
	DIFF	52	46	69	56	55	48
	PROB	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***

YEAR	INBRED	HEATPEAK	HEATBL	BL MOIST	% ROOT LODGE	% STALK LODGE	% DROP EARS	MOISTURE	YIELD
OVERALL	G3001	1350	2484					12.4	71.4
	ZS0560	1422	2578					11.6	66.5
	# EXPTS	27	2					37	37
	DIFF	71	94					0.5	4.8
	PROB	0.000***	0.010***					0.079*	0.023**

YEAR	INBRED	WARM GERM	COLD GERM	% LRG ROUND	% LRG FLAT	% MED ROUND	% MED FLAT	% SML ROUND	% SML FLAT
OVERALL	G3001	97.7	96.7			40.0	23.4		
	ZS0560	97.0	94.0			47.8	16.7		
	# EXPTS	21	21			11	11		
	DIFF	0.7	2.8			7.8	6.7		
	PROB	0.055*	0.002***			0.001***	0.001***		

* .05 < PROB <= .10

** .01 < PROB <= .05

*** .00 < PROB <= .01

Table 3 b shows the inbred G3001 in hybrid combination with a common inbred and another hybrid combination that includes the same common inbred and another inbred. When in these hybrid combinations the present inbred carries less strength in field agronomic traits like lodging and ear drop then the comparison hybrid. However the G3001 hybrid combination brings distinctively significant reduction in moisture levels into the hybrid. The Y/M for the hybrid combination containing the present invention is significantly higher than the comparison hybrid.

Table 4 shows the GCA (general combining ability) estimates of G3001 compared with the GCA estimates of the other inbreds. The estimates show the general combining ability is weighted by the number of experiment/location combinations in which the specific hybrid combination occurs. The interpretation of the data for all traits is that a positive comparison is a practical advantage. A negative comparison is a practical disadvantage. The general combining ability of an inbred is clearly evidenced by the results of the general combining ability estimates. This data compares the inbred parent in a number of hybrid combinations to a group of "checks". The check data is from other companies' hybrids, particularly the leader in the industry and Garst Seed Company's commercial products and pre-commercial hybrids, which were grown in the same sets and locations.

TABLE 4

G3001

	N	FI	Y/M	GI	YLD	MST	%SL	%RL	%DE	TWT	POP	RM
XT=	26	1.1	0.5	0.1	3.8	0.4	-0.8	-1.1	-0.0	-1.2	-1	107

ZS0560

	N	FI	Y/M	GI	YLD	MST	%SL	%RL	%DE	TWT	POP	RM
XT =	60	0.2	-0.2	1.3	1.6	-0.5	0.5	0.0	0.0	-0.1	-209	109

FI= $100 + 0.5(\text{Yld}) - 2.3(\text{MST}) - 0.9(\% \text{SL}) - 0.9(\% \text{RL}) - 2.7(\% \text{DE})$

POP=plants per acre

RM=The Minnesota Relative Maturity

XT= GCA Estimate weighted by parent 2, but using only those parent 2 with two years of data

Table 4 shows G3001 in XT crossed to different inbreds to form 26 hybrid combinations. G3001 in hybrid combination shows an excellent advantage for moisture (MST) and an excellent advantage for yield compared to the commercial checks and the company's commercial inbreds. G3001 has a positive rating for yieldover moisture. In a number of categories the present invention surpasses the ZS0560 line. The ZS0560 line yield advantage is half of the present invention.

TABLE 5A

YIELD RESPONSE

Research Plots

<u>HYBRID</u>	<u>YIELD</u>					
G3001/ Inbred X	66	94	123	152	180	209
Environment	75	100	125	150	175	200

Error: 12.7

#Plots 180

TABLE 5B
YIELD RESPONSE

Research Plots

	<u>HYBRID</u>	<u>YIELD</u>					
5	InbredX/inbred	71	96	120	145	169	194
	Environment	75	100	125	150	175	200

Error: 14.1

#Plots 2700

10

Table 5A shows the yield response of G3001 in hybrid combination in comparison with the plants in the environment around it at the same location. The data for the present inbred is showing consistently better results than the data of the comparison hybrid except in the lowest yielding environments. G3001 in hybrid combination, works well by providing yields better than the environment in high yielding environments. The yield is about equal to yields in the medium yielding environments. Its performance shows that this inbred is best suited to good environments.

15

20

Table 5b shows the data from a different hybrid that was formed with the same inbred X. This comparison hybrid is no longer yielding well particularly in the low and mid environments.

25

The present inbred also carries nice package of average to above average resistance levels to Eyespot, northern leaf blight and other diseases. This resistance allows the yield potential to be met more frequently and in more environments than many other inbreds.

The foregoing is set forth by way of example and is not intended to limit the

scope of the invention.

This invention also is directed to methods for producing a corn plant by crossing a first parent corn plant with a second parent corn plant wherein the first or second parent corn plant is an inbred corn plant from the line G3001. Further, both first and second parent corn plants can come from the inbred corn line G3001. A variety of breeding methods can be selected depending on the mode of reproduction, the trait, the condition of the germplasm. Thus, any such methods using the inbred corn line G3001 are part of this invention: selfing, backcrosses, hybrid production, crosses to populations, haploid by such old and known methods of using stock material that induces haploids and anther culturing and the like. Additionally, this maize can, within the scope of the invention, contain: a mutant gene such as but not limited to sugary 1 or shrunken 1 or waxy or AE or imazethapyr tolerant (IT or IR TM) mutant gene; or transgenic genes such as but not limited to insect resistant genes such as *Bacillus thuringiensis* (Cry genes), or herbicide resistant genes such as Pat gene or Bar gene, EPSP, or disease resistant genes such as the Mosaic virus resistant gene, etc., or trait altering genes such as flowering genes, oil modifying genes, senescence genes and the like.

Various culturing techniques known to those skilled in the art, such as haploid, (stock six is a method that has been in use for twenty years and is well known to those with skill in the art), transformation, and a host of other conventional and unconventional methods are within the scope of the invention. All plants and plant cells produced using the inbred corn line are within the scope of this invention. The term transgenic plant refers to plants having genetic sequences, which are introduced into the genome of a plant by a transformation method and the progeny thereof.

Transformation Methods - are means for integrating new genetic coding

sequences into the plant's genome by the incorporation of these sequences into a plant through man's assistance.

5 Though there are a large number of known methods to transform plants, certain types of plants are more amenable to transformation than are others. Tobacco is a readily transformable plant. The basic steps of transforming plants including monocots are known in the art. These steps are concisely outlined in U.S. patent number 5,484,956 "Fertile Transgenic *Zea mays* Plants Comprising Heterologous DNA Encoding *Bacillus Thuringiensis* Endotoxin" issued January 16, 10 1996 and U.S. patent number 5,489,520 "Process of Producing Fertile *Zea mays* Plants and Progeny Comprising a Gene Encoding Phosphinothricin Acetyl Transferase" issued February 6, 1996.

15 Plant cells such as maize can be transformed by a number of different techniques. Some of these techniques which have been reported on and are known in the art include maize pollen transformation (See University of Toledo 1993 U.S. Patent No. 5,177,010); Biolistic gun technology (See U.S. patent number 5,484,956); Whiskers technology (See U.S. patent numbers 5,464,765 and 5,302,523); Electroporation; PEG on Maize; *Agrobacterium* (See 1996 article on transformation of maize cells in Nature Biotechnology, Volume 14, June 1996) 20 along with numerous other methods which may have slightly lower efficiency rates than those listed. Some of these methods require specific types of cells and other methods can be practiced on any number of cell types.

25 The use of pollen, cotyledons, meristems and ovum as the target issue can eliminate the need for extensive tissue culture work. However, the present state of the technology does not provide very efficient use of this material.

 Generally, cells derived from meristematic tissue are useful. Zygotic embryos can also be used. Additionally, the method of transformation of

meristematic cells of cereal is also taught in the PCT application WO96/04392. Any of the various cell lines, tissues, plants and plant parts can and have been transformed by those having knowledge in the art. Methods of preparing callus from various plants are well known in the art and specific methods are detailed in patents and references used by those skilled in the art. Cultures can be initiated from most of the above-identified tissue. The only true requirement of the transforming material is that it can form a transformed plant. The transgenic gene can come from various non-plant genes (such as; bacteria, yeast, animals, and viruses) along with being from animal or plants.

The DNA used for transformation of these plants clearly may be circular, linear, and double or single stranded. Usually, the DNA is in the form of a plasmid. The plasmid usually contains regulatory and/or targeting sequences which assists the expression of the gene in the plant. The methods of forming plasmids for transformation are known in the art. Plasmid components can include such items as: leader sequences, transit polypeptides, promoters, terminators, genes, introns, marker genes, etc. The structures of the gene orientations can be sense, antisense, partial antisense, or partial sense: multiple gene copies can be used.

The regulatory promoters employed can be constitutive such as CaMv35S (usually for dicots) and polyubiquitin for monocots or tissue specific promoters such as CAB promoters, etc. The prior art includes but is not limited to octopine synthase, nopaline synthase, CaMv19S, mannopine synthase promoters. These regulatory sequences can be combined with introns, terminators, enhancers, leader sequences and the like in the material used for transformation.

The isolated DNA is then transformed into the plant. Many dicots can easily be transformed with *Agrobacterium*. Some monocots are more difficult to transform. As previously noted, there are a number of useful transformation

processes. The improvements in transformation technology are beginning to eliminate the need to regenerate plants from cells. Since 1986, the transformation of pollen has been published and recently the transformation of plant meristems has been published. The transformation of ovum, pollen, and seedlings meristem greatly reduce the difficulties associated with cell regeneration of different plants or genotypes within a plant can present. Duncan, from at least 1985-1988 produced literature on plant regeneration from callus. Both inbred and hybrid callus have resulted in regenerated plants. Somatic embryogenesis has been performed on various maize tissues, which was once considered unusable for this purpose. The prior art clearly teaches the regeneration of plants from various maize tissues.

The most common method of transformation is referred to as gunning or microprojectile bombardment. This Biolistic process has small gold-coated particles coated with DNA shot into the transformable material. Techniques for gunning DNA into cells, tissue, callus, embryos, and the like are well known in the prior art.

After the transformation of the plant material is complete, the next step is identifying the cells or material, which has been transformed. In some cases, a screenable marker is employed such as the beta-glucuronidase gene of the *uidA* locus of *E. coli*. Then, the transformed cells expressing the colored protein are selected for either regeneration or further use. In many cases, a selectable marker identifies the transformed material. The putatively transformed material is exposed to a toxic agent at varying concentrations. The cells not transformed with the selectable marker, which provides resistance to this toxic agent, die. Cells or tissues containing the resistant selectable marker generally proliferate. It has been noted that although selectable markers protect the cells from some of the toxic affects of the herbicide or antibiotic, the cells may still be slightly effected by the toxic agent by having slower growth rates. If the transformed material was cell

lines then these lines are regenerated into plants. The cells' lines are treated to induce tissue differentiation. Methods of regeneration of cellular maize material are well known in the art since early 1982. European Patent Application, publication 160,390, describes tissue culture of corn, which can be used by those skilled in the art. The plants from the transformation process or the plants resulting from a cross using a transformed line or the progeny of such plants are transgenic plants.

Various techniques known to those skilled in the art, such as haploid, transformation, and a host of other conventional and unconventional methods are within the scope of the invention. All plants and plant cells produced using inbred corn line G3001 are within the scope of this invention. The invention encompasses the inbred corn line used in crosses with other, different, corn inbreds to produce (F1) corn hybrid seeds and hybrid plants. This invention includes cells, which upon growth and differentiation produce corn plants having the physiological and morphological characteristics of the inbred line G3001.

A deposit of at least 2500 seeds of this invention will be maintained by Garst Seed Company, 2369 330th Street, Slater, Iowa 50244. Access to this deposit will be available during the pendency of this application to the Commissioner of Patents and Trademarks and persons determined by the Commissioner to be entitled thereto upon request. All restrictions on availability to the public of such material will be removed upon issuance of a granted patent of this application by depositing at least 2500 seeds of this invention at the American Type Culture Collection. The deposit of at least 2500 seeds will be from the same inbred seed taken from the deposit maintained by Garst Seed Company. The ATCC deposit will be maintained in that depository, which is a public depository, for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced if it becomes nonviable during

that period.

Additional public information on some ZS designations may be available from the PVP office a division of the US government.

5

Accordingly, the present invention has been described with some degree of particularity directed to the preferred embodiment of the present invention. It should be appreciated, though, that the present invention is defined by the following claims construed in light of the prior art so that modifications or changes may be made to the preferred embodiment of the present invention without departing from the inventive concepts contained herein.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206